U.S.S.N. 09/779,427

Filed: February 8, 2001

AMENDMENT AND RESPONSE TO OFFICE ACTION

whereby one offers the bacteria at least one substrate carbon source which is selected from the group consisting of: levulinic acid, salts of levulinic acid, esters of levulinic acid, lactones of levulinic acid, substituted levulinic acid or, as the case may be, its derivatives; 5-hydroxyhexanoic acid, its salts, esters and lactones; 4-hydroxyheptanoic acid, its salts, esters and lactones; 4-hydroxyoctanoic acid, its salts, esters and lactones; their halogenated derivatives as well as their mixtures; one incubates the bacteria for a certain time with the carbon source; and one isolates the poly(hydroxy fatty acid) polymers that have been synthesized by the bacteria, it has been possible for the first time to produce 4HV-containing polyesters starting out from levulinic acid. The chemical structure of levulinic acid is reproduced in the following formula:--

In the Claims

1. (Amended) [Process] A process for the preparation of poly(hydroxy fatty acids) comprising incubating a recombinant organism in a mineral medium under aerobic conditions, expressing [with at least one subunit by means of recombinant bacteria which contain and express] at least one fragment of the gene [of] encoding the poly(hydroxy fatty acid) synthase from *Thiocapsa pfennigii* with a substrate carbon source, wherein the recombinant organism produces a poly(hydroxy fatty acid) and,

the poly (hydroxy fatty acid) is recovered [and which are selected from the group comprising:

Pseudomonas putida GPp104 (pHP1014::E156), Alcaligenes eutrophus PHB 4 (pHP1014::EIS6), Pseudomonas putida GPp104 (pHP1014::B28+) [DSM #9417] and Alcaligenes eutrophus PHB 4 (pHP1014:B28+) [DSM # 9418], whereby the bacteria are

AMENDMENT AND RESPONSE TO OFFICE ACTION

•one offers the bacteria at least one substrate carbon source which is selected from the group consisting of:

levulinic acid, salts of levulinic acid, esters of levulinic acid, lactones of levulinic acid, substituted levulinic acid or, as the case may be, its derivatives; 5-hydroxyhexanoic acid, its salts, esters and lactones; 4-hydroxyoctanoic acid, its salts, esters and lactones; their halogenated derivatives as well as their mixtures;

- •one incubates the bacteria for a certain time with the carbon source; and
- •one isolates the poly(hydroxyl fatty acid) polymers that have been synthesized by the bacteria.]
- 2. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein the bacteria are pre-cultivated in a complex medium.
- [4] 3. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein one also adds to the bacterial culture at least one additional carbon source which promotes growth, whereby the carbon source is selected from the group [comprising:] consisting of

octanoic acid salts, eitric acid esters and citric acid lactones, octanoic acid, octanoic acid salts, octanoic acid esters, octanoic acid lactones, [and] gluconic acid, gluconic acid salts, gluconic acid esters, gluconic acid lactones, [; their salts, esters and lactones:] hexoses, [especially glucose and fructose; as well as their mixtures] and combinations thereof.

AMENDMENT AND RESPONSE TO OFFICE ACTION

- 4. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein the process is carried out in the form of a batch process, a fed-batch process, a two-step process or a continuous flow process.
- 5. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein the poly(hydroxy fatty acid) is obtained in a concentration of approximately 15 to 70% by weight [or, especially approximately 15 to 50% by weight or, preferably, approximately 40% by weight] based on the dry mass of the bacterial cells.
- 6. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein the poly(hydroxy fatty acids) are obtained in the form of copolyesters with at least two [or preferably, three] subunits.
- 7. (Amended) [Process in accordance with] The process of claim 1, [characterized by the feature that] wherein the recombinant bacteria are cultivated at cell densities of up to 100 g of dry cellular mass per liter of bacterial nutrient medium.
- 8. (Amended) [Process in accordance with] <u>The process of [claims] claim 1.</u> [characterized by the feature that] <u>wherein</u> one offers the substrate carbon source in excess.
- 9. (Amended) [Process in accordance with] <u>The process of claim 8.</u> [characterized by the feature that] <u>wherein one [uses] offers</u> the substrate carbon source at a concentration of approximately 0.1 to 5% by weight.
- 10. (Amended) [Process in accordance with] The process of claim 9, [characterized by the feature that] wherein one increases the concentration of the substrate carbon source in the

AMENDMENT AND RESPONSE TO OFFICE ACTION

culture medium in steps, optionally with pre-cultivation in the presence of an additional carbon source which does not serve as a substrate.

- 11. (Amended) [Process in accordance with] The process of claim 10, [characterized by the feature that] wherein, [in each case], one adds approximately 0.5% (weight/volume) of neutralized substrate carbon source after approximately 12 h and 24 h at approximately 27°C to 35°C [or, preferably, at approximately 30°C].
- 12. (Amended) [Process in accordance with] The process of claim 1. [characterized by the feature that] wherein cultivation takes place for approximately 24 h to 96 h [or, especially, for approximately 36 h to 72 h or, preferably, for approximately 48 h to 72 h].
- by the feature that] wherein the recombinant bacteria are cultivated under conditions deficient in an element [of deficiency, preferably under] wherein the element is selected from the group consisting of [conditions of a deficiency of] nitrogen, magnesium or phosphate.
- by the feature that] wherein the harvested recombinant bacteria are broken open [by means of physical and/or chemical and/or biochemical processes] in order to obtain the poly(hydroxy fatty acids) that have been produced [bio-technically].
- 15. (Amended) [Process in accordance with] The process of Claim 14, [characterized by the feature that] wherein the harvested recombinant bacteria are lyophilized and then extracted with an organic solvent, [preferably] selected from the group consisting of

AMENDMENT AND RESPONSE TO OFFICE ACTION

chloroform or methylene chloride, in order to break open the recombinant bacteria and to obtain the poly(hydroxy fatty acids).

- by the feature that] wherein the extracted poly(hydroxy fatty acid) produced is precipitated by introducing a hydrophilic solvent, [especially] selected from the group consisting of water [or] and a lower alcohol, [preferably ethanol,] [and] wherein the product is obtained in essentially pure form by removing the hydrophilic solvent.
- by the feature that] wherein the harvested recombinant bacteria are broken open by means [of] selected from the group consisting of detergents, [and/or] a lytic enzyme cocktail, and a combination thereof [as a result of which] wherein the bacterial cell grana, which contain the poly(hydroxy fatty acid), sediment to the bottom of the bio-reactor and are collected from there in order to be processed further.
- 18. (Amended) [Process in accordance with] The process of claim 17, [characterized by the feature that] wherein the lytic enzyme cocktail contains enzymes which are selected from the group [which comprises:] consisting of

lysozyme; proteases; other hydrolytic enzymes; [as well as their mixtures] and combinations thereof.

Please add the following new claims 37-42.

37. The process of claim 1, wherein the poly(hydroxy fatty acid) is obtained in a

AMENDMENT AND RESPONSE TO OFFICE ACTION

- 38. The process of claim 1, wherein the poly(hydroxy fatty acid) is obtained in a concentration of approximately 40% by weight based on the dry mass of the bacterial cells.
- 39. The process of claim 10, wherein, in each case, one adds approximately 0.5% (weight/volume) of neutralized substrate carbon source after approximately 12 h and 24 h at approximately 30°C.
- 40. The process of claim 1, wherein cultivation takes place for approximately 36 h to 72 h.
- 41. The process of claim 1, wherein cultivation takes place for approximately 48 h to 72 h.
- 42. The process of claim 1 wherein the poly(hydroxy fatty acid) polymer is comprised of one or more monomers selected from the group consisting of :
- (A) 3-hydroxybutyric acid, 3 hydroxyvaleric acid and 4-hydroxy-valeric acid;
- (B) 3-hydroxybutyric acid, 3 hydroxyvaleric acid, 4-hydroxy-valeric acid, 3-hydroxyhexanoic acid and 3-hydroxyoctanoic acid;
- (C) 3-hydroxybutyric acid, 3-hydroxyhexanoic acid, 5-hydroxyhexanoic acid, and 3-hydroxyoctanoic acid;
- (D) 3-hydroxybutyric acid, 3 hydroxyvaleric acid, 3-hydroxyhexanoic acid, 3-hydroxyheptanoic acid, 4-hydroxyheptanoic acid and 3-hydroxyoctanoic acid;
- (E) 3-hydroxybutyric acid, 3 hydroxyhexanoic acid, 3-hydroxy-octanoic acid and 4-hydroxyoctanoic acid;